



This fact sheet is developed with funding support from the Land Protection Fund.

Fact sheets are available from Department of Agriculture and Fisheries (DAF) service centres and our Customer Service Centre (telephone 13 25 23). Check our website at www.biosecurity.qld.gov.au to ensure you have the latest version of this fact sheet. The control methods referred to in this fact sheet should be used in accordance with the restrictions (federal and state legislation, and local government laws) directly or indirectly related to each control method. These restrictions may prevent the use of one or more of the methods referred to, depending on individual circumstances. While every care is taken to ensure the accuracy of this information, DAF does not invite reliance upon it, nor accept responsibility for any loss or damage caused by actions based on it.


Department of Agriculture and Fisheries

Biosecurity Queensland

Restricted invasive animal

Wild dog

Canis familaris



The term wild dog refers collectively to purebred dingoes, dingo hybrids and domestic dogs that have escaped or been deliberately released.

Wild dog control methods include baiting, trapping, shooting, fencing, and the use of guardian animals to protect stock. A planned strategy using a combination of these methods that also considers wild dog behavior will enable effective management.

Legal requirments

The wild dog is a restricted invasive animal under the *Biosecurity Act 2014*. It must not be moved, kept (if a dingo), fed, given away, sold, or released into the environment without a permit. The Act requires everyone to take all reasonable and practical steps to minimise the risks associated with invasive plants and animals under their control. This is called a general biosecurity obligation (GBO). This fact sheet gives examples of how you can meet your GBO.



At a local level, each local government must have a biosecurity plan that covers invasive plants and animals in its area. This plan may include actions to be taken on certain species. Some of these actions may be required under local laws. Contact your local government for more information.

Control


Managing wild dogs

To increase wild dog control effectiveness, it is essential that control programs are coordinated among adjoining properties. This fact sheet provides information and some options for controlling wild dogs.

Queensland research has shown that in some situations wild dogs can quickly re-colonise baited areas due to a number of factors including inconsistent bait programs which do not provide comprehensive wild dog control across the landscape. Such programs may alter the dynamics of wild dog populations in the area. To prevent livestock attacks and enhance wild dog management, it is important for producers to work together using a variety of control methods.

Wild dog ecology and seasonal variations can also influence the likelihood of wild dogs coming into contact with a control tool. The timing of control should consider seasonal variations and the availability of water (where water is restricted) and then target watering points. Many land owners bait using 1080 twice a year to target wild dogs during peaks in activity associated with breeding (March/May) and then again in September/November to

Map 1. Distribution of wild dogs in Queensland



target pups and juveniles. However, baiting and trapping is recommended at all times when wild dogs are active.

Fencing

Property fencing suitable to exclude wild dogs is expensive to build and requires continual maintenance to repair damage caused by fallen timber, fire, floods, feral and domestic animals, as well as vegetation regrowth. However, a properly maintained fence can restrict movement into an area where wild dogs have been controlled.

Electric fences suitable for wild dogs have been developed. Electrifying a fence creates a fear of the fence itself and deters wild dogs from approaching.

For property fencing to be successful, the fence must be maintained in good order and ongoing wild dog control conducted within the protected area to limit livestock impacts.

Fencing is the most effective method of protecting livestock and pets from wild dog attack on small acreage blocks.

A fence can also be a good area to place baits and traps when wild dogs are active.

A key success to trapping wild dogs (using foot-hold traps) depends on the skill of the operator. Visit www.feral.org.au to watch a PestSmart video on best practice techniques for wild dog trapping.





For humane reasons and to prevent escape, poisoning traps with strychnine is recommended to quickly kill captured wild dogs. A properly poisoned trap becomes a lethal device rather than a holding device.

A mixture of dog faeces and urine is a popular lure used by trappers. Attractiveness of lures varies with seasons and locations. No single lure has yet been found that is consistently attractive to all wild dogs and repeated use of one lure can lead to aversion amongst remaining dogs.

Traps are best placed in areas of high wild dog activity (known as leads). Here the wild dog is most likely to find and investigate the decoy/odour.

A wild dog scent post (an area where urine or faeces have been deposited) can be found by walking with a domestic dog on a lead along a known pad. Trap placement in relation to the scent post can be optimised by observing the domestic dog's behaviour as it approaches. Factors to consider are:

- where on the bush it smells
- placement of feet while urinating/defecating/sniffing
- how it approaches and where it scratches in relation to the pad and scent post.

Padded, laminated or offset foot-hold traps, in a well tuned and functioning state are recommended.

Shooting

Shooting is an opportunistic method, mostly used for control of small populations or individual problem animals.

Livestock guardian animals

Livestock guardian animals have been used to protect livestock from predators in Europe, Asia and America. Some producers in Queensland have decreased predation on sheep and goats using this method. The use of trapping and poisoning in conjunction with guardian animals must be well planned and managed to ensure guardian animal safety.

Baiting

Poison baits are the most economic, efficient and effective method of controlling wild dogs, especially in inaccessible or extensive areas. Baits can be laid quickly by hand, from vehicles and from aircraft.



Currently there are three poisons legally available for wild dog control. These are 1080 (sodium fluoroacetate), strychnine and para amino propiophenone (PAPP).

Subject to restrictions, 1080 baits, either manufactured or prepared from fresh meat can only be obtained from authorised persons. PAPP can only be supplied as a manufactured bait. A permit from the Queensland Department of Health is required for land owners to purchase strychnine. Strychnine can be used both in baits and on traps. The use of both 1080 and strychnine require adherence to the associated conditions of supply. The use of poison baits will control some but not all wild dogs. Baits should be used in conjunction with all other control tools and not be relied on as a total control method.

Meat baits are attractive both to wild dogs and a range of non-target species. When using meat baits, they can be strategically positioned as wild dogs' keen sense of smell enables them to find baits intentionally buried in sand or otherwise hidden under bushes or in hollow logs. Meat baits may also be tied to prevent their loss to non-target species.

These meat bait placement techniques help to:

- reduce the risk of poisoning non-target species
- increase wild dog contact, hence receiving a lethal
- minimise bait removal by non-target scavengers
- deter ants (ant-covered baits are believed to be less attractive to wild dogs).

Heavy rain within two weeks of baiting can leach 1080 from baits, but baits may still remain toxic for a considerable time.

Ejectors are a new tool in the delivery of 1080. They require a wild dog or fox to pull the ejector head to be activated. This is done by attaching a lure reward to the ejector head. A capsule of lethal dose 1080 is burst into the wild dog's or the foxes mouth. Ejectors are fixed in one stop and are only able to be activated by foxes and dogs.

Further information

Further information is available from your local government office, or by contacting Biosecurity Queensland on 13 25 23 or visit www.biosecurity.qld.gov.au.